

## **Multidisciplinary treatment of lymphedema experience of our center**

### **Complex decongestive therapy experience of a single center**

Topic: physical therapy - surgery

Costantino Eretta\* MD. PHD, Tridoni Elisa\*\*Fkt., Isaia Serena\*\* Fkt, Pigoni Arianna\*\*Fkt., Azzurra Vatteroni\*\*Fkt., Impieri Laura\*\*Fkt., Claudio Bianchi\* MD.

\* General Surgery Department, S.S lymphatic surgery, San Bartolomeo Hospital, Sarzana, Italy

\*\* SME, medical center, Sarzana Italy.

Keywords: lymphedema, disability, complex decongestive therapy (CDT), complex decongestive physiotherapy, manual lymphatic drainage (MLD)

### **Abstract**

Lymphedema is a chronic and worsening disease due to an abnormal accumulation of liquids, with a high protein content in the interstitial space.

The disease is characterized by an insufficient flow of lymphatic fluid, which manifests as edema, inflammation, and fibrosis, all the way up to the stiffening of the affected tissues.

Because it's a chronic and increasing disease, the treatment is highly complex. The literature shows that the treatment must be multidisciplinary and it is necessary to combine multiple techniques, such as manual lymphatic drainage, mechanical lymphatic drainage, elasto-compressive bandages and other complementary techniques up to surgical treatment in the most advanced forms which are not responsive to physical therapy.

Furthermore, the disease is characterized by episodes of cellulitis, that may lead to infectious complications because the lymphatic function becomes insufficient. A condition of local immunodeficiency is created due to the crucial role that the lymphatic system covers with immune defenses, therefore creates a fertile ground for infections caused by small skin wounds, insect bites, animal scratches, nail fungus, blood draws.

Therefore, particular attention is paid to skin folds and interdigital spaces for which hygiene is necessary using neutral detergents, drying by dabbing, and applying emollient creams for skin hydration. Unfortunately, such practices of prevention and care are often underestimated.

We provide education of the patient on self-care, such as the self-bandage and the correct application of the elasto-compressed stocking. For wrapping the bandage, it is advisable to wear a glove or a special sock.

The multilayer bandages are used in the first decongestant phase while over time the elastic stocking is the best aid for the management of lymphedema.

Our clinic is a referral center for diagnosis, treatment and surgical therapy, where patients come for surgical evaluation. Since 2016 we have systematically collected clinical data and centimeter evaluation of more than 600 cases affected by lymphedema and lipoedema of all stages. They have been treated with complex decongestive therapy for 4 hours a day, 15 consecutive days, subsequently a maintenance of once a week for 3 months, then once a month for 6 months.

Of more than 600 patients treated, only 150 were submitted to surgery (lymphatic venous anastomosis, fasciotomy or liposuction, chylothorax and chyloperitoneal shunt, reconstructive plastic of external genitalia).

Conclusion: complex decongestive physical therapy, when properly performed can stabilize the lymphedema patient situation, reducing the stage and ensuring a good quality of life.

## **Introduction**

Lymphedema is a chronic disease with a progressively worsening evolution and appearances of recurrent complications of acute lymphangitic infection, mostly erysipeloid, responsible for a further rapid increase in the volume and consistency of the edema.

The conservative therapeutic methods of medical-physical rehabilitation adopted in the treatment of lymphedema of the limbs allow excellent results to be obtained whether performed by expert hands following precise treatment protocols.

Only patient refractory to conservative treatment should be referred for surgical treatment, a decision exclusively reserved to an experienced lymphologist.

In recent decades, the literature has shown that the advent of surgical techniques such as microsurgery and minimally invasive techniques, including autologous vascularized lymph node microsurgery (VLNT), lymphatic graft (Lympholymphatic graft), anastomosis lymphatic venous vein (LVA) and the superficial one performed in supermicrosurgery, have allowed positive and constantly prolonged results, but they must be always preceded and followed by medical-physical-rehabilitative therapy.

The purpose of our work is to demonstrate how medical physical rehabilitation therapy is effective in treating lymphedema of the limbs, when is performed in highly experienced centers, so then only a small part of patients need surgery.

Our clinic is a referral center for the diagnosis, conservative treatment and surgical therapy for lymphedema, where only physiotherapists with a diploma of Vodder school can work, and always supervised by an expert lymphologist.

Since 2016 we have systematically collected the clinical data and centimeter evaluation of more than 600 cases affected by lymphedema and lipoedema of all stages, treated with complex decongestive therapy for 4 hours a day, 15 consecutive days, subsequently a maintenance of once a week for 3 months, then once a month for 6 months.

Of more than 600 patients treated, only 150 were submitted to surgery (lymphatic venous anastomosis, fasciotomy or liposuction, chylothorax and chyloperitoneal shunt, reconstructive plastic surgery of the external genitalia).

Conclusion: complex decongestive physical therapy, when properly performed can stabilize the lymphedema patient situation, reducing the stage and ensuring a good quality of life

## Patients and Methods.

Between 2016 and 2022 our center conducted a clinical study involving 600 patients with primary and secondary lymphoedema of the limbs undergoing intensive treatments.(table1)

Inclusion and exclusion criteria:

- CDT, patients with clinical stage II or III of lymphoedema (ISL stage), lipoedema all stages; patients with nyha 3 heart failure were excluded from the study;
- SURGERY, patients who could not be stabilized after 2 cycles of intensive care or patients who had an absolute indication for surgical therapy e.g. post-surgical lymphocele, lymphangioma, thoracic duct anomalies, plastic surgery of the external genitalia; patients with lipoedema were excluded from surgery.

The study included 606 patients with middle age of 54 years:

171 with lipoedema, 113 with lower limbs affection (99 women and 14 men), 58 with duplicated lipoedema of upper limbs (56 women and 2 men).

435 with lymphedemas, 125 primary, 119 lower limbs (73 women and 46 men), 6 upper limbs (2 men and 4 women).

About primary lymphedemas, 80 cases were certified for rare disease with exemption code RGG020 (in our region we have been authorized to issue specific exemptions since 2018), 25 of which underwent genetic testing.

291 with secondary lymphedema, 202 lower limbs (123 women and 79 men) for uterin and prostate cancer with lymphadenectomy, a minor part for melanoma with lymphadenectomy, 89 affected by lymphedema of the upper limbs, 82 women (mainly secondary lymphedema to breast cancer) and 7 men (2 for breast cancer and 5 for melanoma surgery).

Patients were classified according to the ISL consensus document:

- primary lymphedema; upper limbs 6 cases stage II, lower limbs 15 stage I, 92 stage II and 12 stage III.

- secondary lymphedema; upper limb 69 cases stage II and 20 cases stage III.

Lower limb lymphedema 171 cases stage II and 31 cases stage III.

19 patients had other pathologies of the truncular lymphatic system.

Table.1

				ISL STAGE			
				CASES	II STAGE		
LIPOEDEMA	LOWER LIMBS	WOMEN	99	113	113		
		MEN	14				
	LOWER LIMBS DUPLICATED UPPER LIMBS	WOMEN	56	58	58		
		MEN	2				
				171			
					ISL STAGE		
				CASES	I STAGE	II STAGE	III STAGE
LYMPHEDEMA	PRIMARY LYMPHEDEMA	LOWER LIMBS	WOMEN	73	15	92	12
			MEN	46			
		UPPER LIMBS	WOMEN	6	0	6	0
			MEN	0			
	SECONDARY LYMPHEDEMA	LOWER LIMBS	WOMEN	123	0	171	31
			MEN	79			
		UPPER LIMBS	WOMEN	82	0	69	20
			MEN	7			
OTHER				19			
				435			
TOTAL CASES				606			

In our center, patients are subjected to a lymphological examination by a specialist with many years of experience. The diagnostic procedure includes the collection of anamnestic data, a clinical examination with venous and arterial Doppler ultrasound examination of the limbs, complete ultrasound abdominal scan, and in case of suspected cancer pathology tumor markers are prescribed. Each patient is discussed collectively with the members of the staff (physiotherapists with a Vodder School diploma).

All patients underwent lymphoscintigraphic examination of the limbs, but for selected or complex cases we address them to MRI studies.

Indications on skin care and how to prevent possible infectious episodes are always explained.

At the beginning of the session, the therapist marks centimeter measurements of limbs, weight and photographic documentation of the patient.

A suitable diet is given for overweight or obese patients.

Our protocol provides an intensive cycle of 15 days of CDP (complex decongestive therapy), each day includes 1 hour of manual lymphatic drainage according to Vodder, 1 hour of LPG Endermology®, 2 hours of sequential compression therapy and a multilayer bandaging with short-stretch bandages.

For patients with facial lymphedema, the protocol included 1 hour of manual lymphatic drainage and the Linforoll® method for one more hour.

Patients who also presented web axillar syndrome or lymphatic drainage disturbed by scars, we combined the lymph taping.

At the end of the CDP, the patient's measurements are taken again, so the clinical response can be evaluated. The specialist then prescribes the correct elastic stocking with the right compression class, which is supplied by our trusted technician based on accurate measurements, so one or more customized elastic garments are packaged.

Stabilized patients continue with 1 weekly sessions for a further 3 months and subsequently 1 session a month for 6 months, in our center or with a Vodder physiotherapist near the patient's residence.

Patients refractory after at least 2 intensive cycles of CDP may be cases for surgery indications.

In our center, the following surgical procedures are performed based on the residual edema and its component (fibrotic vs adipose) and on the lymphoscintigraphic study:

- One site deep lymphatic-venous anastomosis (LVA) In the inguinocrural region for the lower limb and in the middle third of the arm for the upper limb, using the blue patent violet and operating microscope.
- Supermicrosurgery, with the use of indocyanine green, with multiple lymphatic venous anastomoses.
- MIF, minimally invasive endoscopic fasciotomy for cases where is not possible to perform lymphatic-venous anastomosis or for oncological criteria.
- Fibrolympholiposuction, in patients with a higher percentage of residual adipose tissue.
- Resective surgery in those patients where exuberant tissue remains at the end of the decongestant therapy or in patients presenting asymmetrical lipodystrophy, resective plastic surgery of the external genitalia, removal of cystic lymphangioma, lymphocele.
- Treatment of chylothorax and chyloperitoneum in patients with abnormalities of the thoracic duct or of the chyliferous cistern, who are preventively undergoing hospitalization and total parenteral therapy and subsequently laparoscopic or thoracoscopic video surgery.
- Implantation of monoclonal cells, in patients with lymphedema and ulcerations.

## Results

In our center, we treated all patients with CDP according to our protocol, and in order to evaluate the response to the treatment we decided to use the ICF Disability Scale. The ICF is outlined as a classification that aims to describe the state of people's health in relation to each field of life (social, family, work) in order to highlight the difficulties that can cause disability in the socio-cultural environment of the patient. The ICF Disability Scale is a framework of predefined tables which each patient is asked to fill out at the beginning and at the end of the cycle of treatments.

Thanks to this methodology we have highlighted the physical and psychological difficulties of the patient during the various stages of the disease. The aim was to verify the patient's improvements by comparing the results obtained based on the subjective responses they made.

The questionnaires collected data about the disability degree that they express personally, demonstrating how the proposed treatments were able to improve psycho-physical and social conditions of the patient suffering from lymphedema and lipedema. (tab.2-3)

**Tab 2. ICF UPPER LIMB QUESTIONNAIRE**

SKILL	0	1	2	3	4
LIFT AND CARRY OBJECT					
FINE MOVEMENT OF THE HAND					
VEHICLE					
SELF CLEANING					
CARING OF SINGLE PARTS OF THE BODY					
DRESSING					
PREPARING MEALS					
HOUSEWORKS					
CARING OF HOUSE THINGS					
SIMPLE INTERPERSONAL INTERACTIONS					
COMPLEX INTERPERSONAL INTERACTIONS					
INTERACTING WITH NEW PEOPLE					
FORMAL RELATIONS					
INFORMAL RELATIONS					
FAMILY RELATIONS					
INTIMATE RELATIONS					
JOB					
FREE TIME					
RELIGION AND SPIRITUALITY					
POLITICAL ACTIVITY					

**Tab 3. ICF LOWER LIMB QUESTIONNAIRE**

SKILL	0	1	2	3	4
CHANGE BODY POSITIONS					
KEEP BODY POSITION					
MOVE IN SPACE					
WALKING					
VEHICLE					
DRIVING					
CARING OF SINGLE PARTS OF THE BODY					
BODY NEEDS					
DRESSING					
HOUSEWORKS, CARING OF HOUSE THINGS					
SIMPLE INTERPERSONAL INTERACTIONS					
COMPLEX INTERPERSONAL INTERACTIONS					
INTERACTING WITH NEW PEOPLE					
FORMAL RELATIONS					
INFORMAL RELATIONS					
FAMILY RELATIONS					
INTIMATE RELATIONS					
JOB					
FREE TIME					
RELIGION AND SPIRITUALITY					
POLITICAL ACTIVITY					

In order to get the Disability Average Score, the patient has to express each skill with a score between 0 and 4. Then the professional calculate the total amount by adding each single score and dividing it by the total number of items. The number obtained is related to an index that describe the degree of disability.(tab.4).

*Tab.4 Disability Average Score.*


<b>0</b>	<b>ABSENCE OF DISABILITY 0 – 0,5</b>	Patients do their activity with no restrictions in choosing, or problems in any function, even with orthoses
<b>1</b>	<b>MINOR DISABILITY 0,6 – 1,5</b>	Patients have minor disability in at least one activity
<b>2</b>	<b>MODERATE DISABILITY 1,6 – 2,5</b>	Patients have moderate disability in at least one activity
<b>3</b>	<b>SERIOUS DISABILITY 2,6 – 3,5</b>	Patients have serious disability in at least one activity
<b>4</b>	<b>COMPLETE DISABILITY 3,6 - 4</b>	Patients have complete disability in at least one activity



All patients affected by lipoedema have been treated with CDP, and those with BMI higher than 25 had to follow a suitable diet combined with regular physical activity, getting benefits from both decongestant therapy and weight loss.

All patients reported reduction of pain, volume and sensation of tension (fig.1)

Figura 1. *Patients affected by Lipoedema at lower limb in different steps while making CDP.*



About patients affected by lymphedema of all 435 who underwent CDP, 285 had a good reduction of volume, in some cases up to the complete normalization of the edema and reduction of cellulite episodes. All patients wore elastic stocking in the maintenance phase and still continue to take sessions of Vodder Manual Lymphatic Drainage, depending on the residual clinical stage. (fig.2-3-4-5-6-7-8)

Examples below: patients affected by lymphedema of limbs, before and after CDP.




Fig. 2 *Breast Cancer*




Fig. 3 *Hysterectomy*




Fig. 4 *Prostatectomy*




Fig. 5 *Hysterectomy*




Fig. 6 *Primary Lymphedema*





Fig.7 Primary lymphedema

Fig. 8 Stabilization of lymphedema with elastic stocking



The 135 patients refractory to conservative therapy after two intensive cycles, 6 months apart, underwent surgery. The indication was placed on the basis of lymphoscintigraphy, the stage of the edema and the component of the edema itself (fibrotic/adipose tissue), the localization of the edema or the congenital anomalies, e.g. anomalies of thoracic duct or chylous cistern. (Tab 4).

N° patients	Type of Surgery	Pathology
89	LVA one site	Lymphedema
9	LVA multiple	Lymphedema
2	VLT	Chylothorax
13	Resective surgery	Lymphocele post surgery
1	VLE	Chyloperitoneum
5	Plastic surgery of genitalia	Genitalia Lymphedema
20	Mini-invasive Fasciotomy	Lymphedema
10	Fibrolymphosuction	Lymphedema with fibrotic-adipose tissue
1	Monoclonal cell implantation	Lymphedema with ulcer
Tot 150		

The patients subjected to the different surgical techniques benefited from the result by stabilizing the residual edema and in many cases improving the measurements of volume. Some patients underwent also to a combined technique. Fig. 9-10-11-12-13-14-15-16-17-18.

Fig 9 *LVA one site with microscope*




Fig.10 *Secondary lymphedema result one site LVA + CDT*






Fig.11 *Secondary lymphedema result multiple LVA + CDT*




Fig.12 *MIF (mini-invasive fasciotomy)*




Fig.13 *Primary lymphedema MIF result*




Fig.14 *Plastic surgery of genitalia*






Fig.15 *Resective Surgery*




Fig.16 *Fibrolymphosuction technique with PDE*




Fig.17 *Fibrolymphosuction with CDP*





Fig. 18 *Monoclonal cell implantation in primary lymphedema with ulcer, with PDE*




The results collected demonstrate that when the combined complex decongestant therapy is conducted in specialized referral centres, with a qualified team of professionals in collaboration with an expert lymphologist, that is profoundly effective to control the lymphedema and lipoedema process in the majority of cases (less than 20 percent of patients need for surgical therapy).

Based on the data obtained using the ICF Score, the subjects under examination were able to express themselves, demonstrating the improvement of the psycho-physical and social conditions of their lives, feeling great satisfaction with the treatments received. (tab. 5-6)


We integrated the reading of the results also with the radar charts, for a better consultation and understanding, especially for the patients, so they can easily recognize the changes.

In this way it's clear to see in a unique image how the general situation before the treatment located mostly in the area of moderate disability, than after the treatment moved to a new area between minor and absence of disability, thanks to the changes of the data.

Tab 5. Results for ICF lipoedema



Tab 6. Results for ICF lymphedema



## Conclusions

Lymphedema is a disease which is very difficult to manage for patients, mostly because of the disability that the disease manifest in their life. At the most evident level there are the physical changes of the body, especially with limbs that became swollen, stiff, painful and easy to get infections and complications due to the immunological impairment of a decreased lymph flow.

But beyond the physical manifestations there is a huge impact on the psychological level, in private and social life.

Our study has finally demonstrated that a well conducted strategy of therapies, known as Complex Decongestive Therapy (CDP) performed by specialized professionals can lead to astounding results for the patients, in terms of both data and disabilities.

A proper Decongestive Therapy is made first of all by the Manual Lymphatic Drainage that only a Vodder professional can execute in a precise way, thanks to the deep knowledge and sensations that is reached by the therapist in order to adapt each time to the different tissue that the hand find. This can make a great difference.

Vodder Manual Lymphatic Drainage associated with the right compression of bandages and stocking in following stages are the real CDP that a patient has to find in a serious center.

The real CDP has a fantastic numbers of demonstrations in literature about effectiveness and tangible improvement of the quality of life found by people affected by lymphedema or lipoedema disease.

## References

1. Becker C,Arrive L,Saaristo A,Germain ,Fanzio P,Batista BN, Piquilloud G,  
*Surgical treatment of congenital lymphedema.*Clin Plast Surg.2012 Oct;39(4):377-84.
2. Benda.K ,Lebloch D.,Bendova.M. *Prevention of primary lymphedema-possible way,*Lymphology,31 (suppl) 1998:465-468.
3. Bernard P. *Primary and secondary hospitalization criteria.* Ann Dermatol Venereol. 2001 Mar; 128(3 Pt 2):363-7.
4. Boccardo.F, Campisi C et al. *A Pilot. Study of prevention of secondary lymphedema.*Lymphology 2000;33:222-555.
5. Boccardo F, Bellini C, Eretta C, Pertile D, Da Rin E, Benatti E, Campisi M, Talamo G, Macciò A, Campisi C, Bonioli E, Campisi C. *The lymphatics in the pathophysiology of thoracic and abdominal surgical pathology: immunological consequences and the unexpected role of microsurgery.* Microsurgery. 2007;
6. Br J Community Nurs. *Improving patient concordance in lymphoedema management with SoftFit techonology .*2017 May.
7. Bruna J:*Indication for lymphography in the era of new imaging methods.* Lymphology 1994,27 (Suppl),319-320.
8. Bonnetblanc JM, Bedane C. *Erysipelas: recognition and management.* Am J Clin Dermatol. 2003; 4(3): 157-63.
9. B.A.Corliss,M.S.Azimi,J.Munson,S.M.Peirce,W.L.Murfee “*Macrophages: An inflammatory link between Angiogenesis and Lymphangiogenesis*”, microcirculation 2016 february.
10. Campisi C, Bellini C, Eretta C, Zilli A, da Rin E, Davini D, Bonioli E, Boccardo F. *Diagnosis and management of primary chylous ascites.* J Vasc Surg. 2006 Jun;43(6).
11. Campisi C, Davini D, Bellini C, Taddei G, Villa G, Fulcheri E, Zilli A, Da Rin E, Eretta C, Boccardo F. *Lymphatic microsurgery for the treatment of lymphedema.* Microsurgery. 2006;
12. Campisi C, Davini D, Bellini C, Taddei G, Villa G, Fulcheri E, Zilli A, da Rin E, Eretta C, Boccardo F. *Is there a role for microsurgery in the prevention of arm lymphedema secondary to breast cancer treatment?* Microsurgery. 2006;
13. Campisi C, Eretta C, Pertile D, Da Rin E, Campisi C, Macciò A, Campisi M, Accogli S, Bellini C, Bonioli E, Boccardo F. *Microsurgery for treatment of peripheral lymphedema: long-term outcome and future perspectives.* Microsurgery. 2007;27(4):333-8

14. Campisi C, Zilli A, Macciò A, Schenone F, da Rin E, Eretta C, Boccardo F. *La prevenzione del linfedema secondario al trattamento del tumore della mammella: dal caso clinico ad una proposta di protocollo di prevenzione*. Chir Ital. 2004 May-Jun;56(3):419-24. Italian
15. Cooper G: *compression of therapy and the management of lower-limb lymphoedema: the male perspective*. Br J Community Nurs. 2015 Mar;20(3):118,120,122-4.
16. De Godoy JM, De Godoy MF, Valente A, Camacho EL, Paiva EV. *Lymphoscintigraphic evaluation in patients after erysipelas*. Lymphology 2000 Dec;
17. Dupuy A, Benchikhi H, Roujeau JC, Bernard P, Vaillant L, Chosidow O, Sassolas B, Guillaume JC, Grob JJ, Bastuji-Garin S. *Risk factors for erysipelas of the leg (cellulitis): case-control study*. BMJ 1999 Jun 12;
18. Eretta C, Ferrarese A, Moggia E, Francone E, Sagnelli C, Martino MD, de Franciscis S, Amato B, Grande R, Butrico L, Amato M, Serra R, Martino V, Berti S. *Surgical treatment of recidivist lymphedema*. Open Med (Wars). 2016 Jun 23;
19. Executive Committee of the international Society of Lymphology, *The diagnosis and treatment of peripheral lymphedema: 2020 Consensus Document of the International Society of Lymphology*. Lymphology. 2020;53(1):3-19.
20. Földi E. *Therapy of lymphedema*. Hautarzt. 2012 August;63(8):627-33.
21. Földi M. *The therapy of lymphedema*. European Journal of Lymphology and related problems 1993-1994;14:43-49,59;
22. G. Bonetti, K. Dhuli, S. Michelini, S. Michelini, S. Michelini, M. Ricci, M. Cestari, M. Bertelli, *Dietary supplements in lymphedema*, 2022 Oct 17;63.
23. Honnor A1: *Understanding the management of lymphoedema for patients with advanced disease*. Int J Palliat Nurs. 2009 Apr;15(4):162,164,166-9.
24. International Society of Lymphology Executive Committee. *The Diagnosis and Treatment of Peripheral Lymphedema*. Lymphology 28 (1995).
25. International Congress of Lymphology, Chennai, India. General Assembly ISL *Consensus Document Revisited*. September 25, 1999.
26. ISL Executive Committee Meeting, Földi Klinik, Hinterzarten, Germany. Discussions on modification of the ISL *Consensus Document*. August 30.
27. Ishida O et Al.: *Evaluation of lymphatic and non lymphatic edema by MRI*. in progress in Lymphology XIII, Ed. EV Cluzan, Elsevier Sc. Publisher, 1992.



28. Kano Y, Inaoka M, Shiohara T. *Superficial lymphangitis with interface dermatitis occurring shortly after a minor injury: possible involvement of a bacterial infection and contact allergens.* Dermatology 2001; 203(3):217-20.
29. Lasinski BB; *Complete decongestive therapy for treatment of lymphedema.* Semin Oncol Nurs 2013 Feb;29(1):20-7.
30. Leduc A; *Le drainage lymphatique.* 7th ed, Parigi, Masson, 47-55, 1991
31. Leduc A, Bourgeois P, Bastin R; *Lymphatic reabsorption of proteins and pressotherapies.* V Congrès du Group Européen de Lymphologie (GEL) Porto, 1985
32. M.L.Lanzi , *Pedagogia Sociale Manuale per l'infermiere*, Carocci Faber, 2004.
33. Mancini S; *Trattato di Flebologia e Linfologia*, UTET, 2001.
34. Michelini S, Caldirola R, Forner Cordero I, Olszewski WL, Pissas A, Dimakakos E, Michelotti L. *"Linforoll: A new device for treatment of lymphedema. Preliminary experience"* Eur J Lymph 2013;24,25
35. Michelini S, Paolacci S, Manara E, Eretta C, Mattassi R, Lee BB, Bertelli M. *Genetic tests in lymphatic vascular malformations and lymphedema.* J Med Genet. 2018 Apr.
36. Mihara M, Hara H, Kikuchi, Yamamoto T, Lida T, Narushima M, Araky J, Murai N, Mitsui K, Gennaro P, Gabriele G, Koshima I, *Scarless lymphatic venous anastomosis for latent and early-stage lymphoedema using indocyanine green lymphography and non-invasive instruments for visualising subcutaneous vein.* J Plast Reconstr Aesthet Surg. 2012 Nov;65 (11).
37. Mihara M, Murai N, Hayashi Y, Hara H, Lida T, Narushima M, Todokoro T, Uchida G, Yamamoto T, Koshima I, *Using indocyanine green fluorescent lymphography and lymphatic venous anastomosis for cancer related lymphedema.* Ann Vasc Surg. 2012 Feb;26(2):278.e1-6.
38. *The Diagnosis and Treatment of Peripheral Lymphedema: 2009 Consensus Document of the International Society of Lymphology.* Lymphology 42 (2009).
39. *The Diagnosis and Treatment of Peripheral Lymphedema: 2013 Consensus Document of the International Society of Lymphology.* Lymphology 46 (2013).
40. Thomson M, Walker J.; *Collaborative lymphoedema management: developing a clinical;* Int J Palliat Nurs, 2011 May;17(5):231-8.
41. Uhara H, Saida T, Watanabe T, Takizawa Y. *Lymphangitis of the foot demonstrating lymphatic drainage pathways from the sole.* J Am Acad Dermatol. 2002 Oct;47(4):502-4.
42. Vaillant L. *Diagnostic criteria for erysipelas.* Ann Dermatol Venereol. 2001 Mar; 128 (3 Pt 2): 326-33.